

# Lower Mass Limit for the Axion

Kirk T. McDonald

*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*

(May 7, 2026)

The axion is a (pseudoscalar) spin-0 boson predicted by Wilczek [1] and Weinberg [2] as a solution to the so-called strong- $CP$  problem identified by Peccei and Quinn [3].

Since axions are bosons they can in principle form a Bose-Einstein condensate with characteristic radius  $L$ . There is recent evidence [4] that the formation of such a condensate has a coherence length  $l \leq L$  whose growth is related by

$$\frac{dl^2}{t} = \frac{3.4\hbar}{m}, \quad (1)$$

where  $m$  is the mass of the boson. Supposing that  $dl/dt$  is limited by the speed of light,  $c$ , we infer that

$$m \geq \frac{1.7\hbar}{cl} \approx \frac{6 \times 10^{-11} m_e}{l}, \quad (2)$$

for  $l$  in meters, where  $m_e \approx 10^{-30}$  kg is the mass of the electron. For  $l \approx L = 1$  m, as might be achieved in an experiment [5] where axions are confined in a long optical fiber coiled into a circle of 1-m diameter inside a strong magnetic field, eq. (2) implies that  $m \gtrsim 6 \times 10^{-11} m_e$  and  $m/c^2 \gtrsim 3 \times 10^{-5}$  eV which is somewhat above the lower end of the stated range of sensitivity of the experiment. For a fiber coiled into a circle of 10-cm diameter the mass limit would be 10 times higher.

## References

- [1] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, Phys. Rev. Lett. **40**, 279-282 (1977). <https://doi.org/10.1103/PhysRevLett.40.279>  
[https://kirkmcd.princeton.edu/examples/EP/wilczek\\_prl\\_40\\_279\\_78.pdf](https://kirkmcd.princeton.edu/examples/EP/wilczek_prl_40_279_78.pdf)
- [2] S. Weinberg, *A New Light Boson?* Phys. Rev. Lett. **40**, 223-226 (1978).  
<https://doi.org/10.1103/PhysRevLett.40.223>  
[https://kirkmcd.princeton.edu/examples/EP/weinberg\\_prl\\_40\\_223\\_78.pdf](https://kirkmcd.princeton.edu/examples/EP/weinberg_prl_40_223_78.pdf)
- [3] R.D. Peccei and H.R. Quinn, *CP Conservation in the Presence of Pseudoparticles*, Phys. Rev. Lett. **38**, 1440-1443 (1977). <https://doi.org/10.1103/PhysRevLett.38.1440>  
[https://kirkmcd.princeton.edu/examples/EP/peccei\\_prl\\_38\\_1440\\_77.pdf](https://kirkmcd.princeton.edu/examples/EP/peccei_prl_38_1440_77.pdf)
- [4] G. Martirosoyan *et al.*, *A universal speed limit for spreading of coherence*, Nature **647**, 608-612 (2025). <https://doi.org/10.1038/s41586-025-09735-z>  
[https://kirkmcd.princeton.edu/examples/QM/martirosoyan\\_nature\\_647\\_608\\_25.pdf](https://kirkmcd.princeton.edu/examples/QM/martirosoyan_nature_647_608_25.pdf)
- [5] D.D. Yavuz and S. Inbar, *Generation and detection of axions using guided structures*, Phys. Rev. D **647**, 074012 (2022). <https://doi.org/10.1103/PhysRevD.105.074012>  
[https://kirkmcd.princeton.edu/examples/EP/yuvaz\\_prd\\_105\\_074012\\_22.pdf](https://kirkmcd.princeton.edu/examples/EP/yuvaz_prd_105_074012_22.pdf)